



Contribution ID: 90 Contribution code: FRBO4

Type: Contributed Oral

A Perfect X-Ray Beam Splitter and its Applications to Time-Domain Interferometry and Quantum Optics Exploiting FELs

Friday, 26 August 2022 12:25 (25 minutes)

Brilliant, ultrashort, and coherent X-ray FEL pulses allow investigations of dynamics at the inherent time and length scale of atoms. However, the user community still lacks access to phase-locked X-ray pulses, desirable for time domain correlation spectroscopies and coherent quantum control. Based on selective electron-bunch degradation in the accelerator, combined with two-stage, self-seeded photon emission, we propose an FEL mode generating subfemtosecond, phase-locked X-ray pulse pairs with up to 100 fs delay. Splitting the electron bunch in the accelerator, instead of photon pulses in the beamline, avoids relative phase jitter. This enables time-domain interferometry, such as the X-ray analog of the ubiquitous Fourier transform infrared spectrometer, and, more generally, all of nonlinear and quantum optics requiring coherent copies of beams.

I have read and accept the Privacy Policy Statement

Yes

Primary authors: REICHE, Sven (Paul Scherrer Institut); PEDRINI, Bill (Paul Scherrer Institut); PRAT, Edouard (Paul Scherrer Institut); AEPPLI, Gabriel (Paul Scherrer Institut and ETH Zurich); Dr GERBER, Simon (Paul Scherrer Institut)

Presenter: REICHE, Sven (Paul Scherrer Institut)

Session Classification: End-to-end experiments (machine driven)

Track Classification: End-to-end experiments (machine driven)