



Contribution ID: 164 Contribution code: WEP05

Type: Contributed Poster

## Wakefield Calculations of the Undulator Section in FEL-I at the SHINE

*Wednesday, 24 August 2022 17:10 (20 minutes)*

In free electron lasers (FEL) the accumulative effects of wakefields always lead to critical impacts on the electron bunch, resulting in an energy spread and deviation of transverse position. Thus the lasing performance will be decreased. The Shanghai high-repetition-rate XFEL and extreme light facility (SHINE) is under construction and the wakefields estimations are required. The SHINE contains three different undulator lines (FEL-I- III) designed for different functions. The wakefields of FEL-I undulator section has been studied in our work before. However the wakefields of inner segments between undulators are calculated simply. In this paper, we calculate the wakefields of inner segments considering more exquisite structures in FEL-I. We consider gradual changed connections between beam pipes of different diameters and dechirpers. We compared wakefields of different schemes of inner segments. Based on the results, we give some suggestions for the designation of the inner segments in FEL-I.

### I have read and accept the Privacy Policy Statement

Yes

**Primary authors:** LIU, He (Shanghai Institute of Applied Physics); YAN, Jiawei (European XFEL GmbH); GUO, Jun-Jie (Zhangjiang Laboratory Shanghai); LIU, Tao (Shanghai Advanced Research Institute Chinese Academy of Science); DENG, Haixiao (Shanghai Advanced Research Institute Chinese Academy of Science); LIU, Bo (Shanghai Advanced Research Institute Chinese Academy of Science)

**Presenters:** LIU, He (Shanghai Institute of Applied Physics); YAN, Jiawei (European XFEL GmbH); DENG, Haixiao (Shanghai Advanced Research Institute Chinese Academy of Science); LIU, Bo (Shanghai Advanced Research Institute Chinese Academy of Science)

**Session Classification:** Wednesday posters

**Track Classification:** Electron beam dynamics