



Contribution ID: 69 Contribution code: MOP55

Type: Contributed Poster

## Superradiant Amplification to Produce Attosecond Pulses in Soft X-Ray Regime via Linear Reverse Taper within Undulator Section

*Monday, 22 August 2022 17:10 (20 minutes)*

Laser pulses of sub-femtosecond duration can be used to track the motion of electrons in the inner shell, which is needed in a variety of advanced experiments. Although this has been accomplished in XUV and hard X-rays in a free-electron-laser facility, it remains a challenge in the soft X-ray region due to the relatively high photon energies and large slippage in the undulator. In this contribution, we present a method to achieve a pulse sequence of  $\sim 120$  attosecond each in average at 293.8 eV photon energy (4 nm wavelength), which covers the K-shell absorption of Carbon. The key is to create a linear undulator taper within each undulator module by rotating a transverse gradient undulator (TGU) at a small angle. The TGU technique is usually referred to minimise the energy spread effect in Laser-driven plasma accelerator, while in this paper we demonstrate that it can also be used to generate short pulses.

### I have read and accept the Privacy Policy Statement

Yes

**Primary author:** Mr ZHU, Longdi (Paul Scherrer Institut)**Co-authors:** REICHE, Sven (Paul Scherrer Institut); FERRARI, Eugenio (Deutsches Elektronen-Synchrotron)**Presenters:** Mr ZHU, Longdi (Paul Scherrer Institut); REICHE, Sven (Paul Scherrer Institut); FERRARI, Eugenio (Deutsches Elektronen-Synchrotron)**Session Classification:** Monday posters**Track Classification:** SASE FEL