



Contribution ID: 146 Contribution code: MOCO3

Type: **Contributed Oral**

Proposal for a Quantum Free Electron Laser Driven by Ultracold Electrons

Monday, 22 August 2022 15:10 (25 minutes)

Operation of a Quantum Free-Electron Laser (QFEL) could provide fully coherent X- and gamma-rays in a compact setup. Imperative to experimental realization is allowing for decoherence of either spontaneous emission or space-charge to take place, having opposing constraints [1]. Here, for the first time, we discuss a comprehensive QFEL model that takes into account both decoherence effects. Then, we use this model to investigate the ultracold electron source (UCES) [2] as a potential QFEL electron injector. The UCES, based on near-threshold photoionization of laser-cooled and trapped atomic gas, has the unique property of allowing highly charged electron bunches to be extracted while maintaining ultralow transverse emittance. We find that the ultracold electron bunches meet the stringent requirement for potential QFEL operation with commercially available laser systems.

I have read and accept the Privacy Policy Statement

Yes

Primary authors: SCHAAP, Brian (Technische Universiteit Eindhoven); LUITEN, Jom (Technische Universiteit Eindhoven); Mr SCHOUWENAARS, Sander (Technische Universiteit Eindhoven)

Presenters: SCHAAP, Brian (Technische Universiteit Eindhoven); LUITEN, Jom (Technische Universiteit Eindhoven); Mr SCHOUWENAARS, Sander (Technische Universiteit Eindhoven)

Session Classification: FEL Theory

Track Classification: FEL Theory