



Contribution ID: 1069 Contribution code: MOPG09

Type: **Poster Presentation**

A faster algorithm to compute lowest order longitudinal and transverse resistive wall wake for non-ultrarelativistic case

Monday, 20 May 2024 16:00 (2 hours)

With the development of the steady state micro bunching (SSMB) storage ring, its parameters reveal that the ultra relativistic assumption which is wildly used is not valid for the electron beam bunch train, which has length in the 100 nm range, spacing of 1 μm and energy in hundreds MeV range. The strength of the interaction between such bunches and the potential instability may need careful evaluation. At the same time, the effect of the space charge inside a single bunch due to space charge effect also needs to be considered. In this article, we reorganized the lowest-order longitudinal wakefield under non-ultra relativistic conditions, and modified the inconsistent part in the theoretical derivation in some essays of the lowest-order transverse wakefield. We present the modified theoretical results and analysis. Then based on the result we have derived, we give a algorithm which is thousands time faster than direct calculation. It lays foundation in future research.

Footnotes

Funding Agency

Paper preparation format

LaTeX

Region represented

Asia

Primary author: Mr TANG, Jiazhen (Tsinghua University in Beijing)

Presenter: Mr TANG, Jiazhen (Tsinghua University in Beijing)

Session Classification: Monday Poster Session

Track Classification: MC2: Photon Sources and Electron Accelerators: MC2.A04 Circular Accelerators