



Contribution ID: 1609 Contribution code: THPC21

Type: Poster Presentation

## Measured dynamic aperture and detuning of nonlinear integrable optics

*Thursday, 23 May 2024 16:00 (2 hours)*

One of the most promising advantages of nonlinear integrable optics is strong amplitude dependent tune shift without degrading the dynamic aperture. The integrable optics test accelerator (IOTA) at Fermilab is constructed around nonlinear lattice elements of the elliptical type as described by Danilov and Nagaitsev. Detuning and dynamic aperture scans in IOTA were performed using a fast dipole kicker and a low emittance electron beam. The evolution of the dynamic aperture and detuning for different configurations of the integrable optics lattice are presented.

### Footnotes

### Funding Agency

### Paper preparation format

LaTeX

### Region represented

North America

**Primary author:** WIELAND, John (Fermi National Accelerator Laboratory)

**Co-authors:** ROMANOV, Alexander (Fermi National Accelerator Laboratory); VALISHEV, Alexander (Fermi National Accelerator Laboratory); KUKLEV, Nikita (Argonne National Laboratory)

**Presenter:** WIELAND, John (Fermi National Accelerator Laboratory)

**Session Classification:** Thursday Poster Session

**Track Classification:** MC5: Beam Dynamics and EM Fields: MC5.D01 Beam Optics Lattices, Correction Schemes, Transport