



Contribution ID: 2074 Contribution code: SUPC004

Type: Student Poster Presentation

## Numerical methods for emittance computation from luminosity

*Sunday, 19 May 2024 14:00 (4 hours)*

The beam transverse emittances play a critical role in high-energy colliders. Various measurement techniques are employed to measure them. In particular, the so-called luminosity emittance scans (or Van der Meer scans) are used in order to evaluate the convoluted beam emittances. This method assumes different emittances in the two planes but identical emittances in the two beams. In this paper, we propose an approach to remove this constraint. After having presented the new measurement protocol, we will discuss its potential and limits, including the statistical measurement error of the luminosity value as obtained from numerical studies.

### Footnotes

### Funding Agency

### Paper preparation format

### Region represented

Europe

**Primary author:** RUFOLO, Matteo (IDSIA Dalla Molle Institute for Artificial Intelligence USI-SUPSI)

**Co-authors:** FORNARA, Andrea (The University of Manchester); LAMB, Eleanor (Ecole Polytechnique Fédérale de Lausanne); STERBINI, Guido (European Organization for Nuclear Research); GIACOMEL, Lorenzo (European Organization for Nuclear Research)

**Presenter:** RUFOLO, Matteo (IDSIA Dalla Molle Institute for Artificial Intelligence USI-SUPSI)

**Session Classification:** Student Poster Session

**Track Classification:** MC1: Colliders and other Particle and Nuclear and Physics Accelerators: MC1.A01 Hadron Colliders