



Contribution ID: 2130 Contribution code: SUPC027

Type: Student Poster Presentation

Various methods for computing dominant spin-orbit resonance strengths in storage rings

Sunday, 19 May 2024 14:00 (4 hours)

The strength of a first-order spin-orbit resonance is defined as the amplitude of the corresponding Fourier component of the spin-precession vector. However, it is possible to obtain the resonance strength without computing the Fourier integral directly. If a resonance is sufficiently strong, then to a good approximation, one can neglect all other depolarizing effects when near the resonance. Such an approximation leads to the single resonance model (SRM), for which many aspects of spin motion are analytically solvable. In this paper, we calculate the strength of first-order resonances using various formulae derived from the SRM, utilizing spin tracking data, the direction of the invariant spin field, and jumps in the amplitude-dependent spin tune. Examples are drawn from the RHIC Blue ring.

Footnotes

Funding Agency

Supported by DOE under No. DEAC0298-CH10886, DE SC-0024287, and DE-SC0018008.

Paper preparation format

LaTeX

Region represented

North America

Primary author: DEVLIN, Joseph (Cornell University (CLASSE))

Co-author: HOFFSTAETTER, Georg (Cornell University (CLASSE))

Presenter: DEVLIN, Joseph (Cornell University (CLASSE))

Session Classification: Student Poster Session

Track Classification: MC1: Colliders and other Particle and Nuclear and Physics Accelerators: MC1.A24 Accelerators and Storage Rings, Other