



Contribution ID: 2179 Contribution code: SUPC082

Type: **Poster Presentation**

## An ultimate single-ion source using a Coulomb crystal in a Paul trap

Sunday, 19 May 2024 16:00 (2 hours)

An ion cloud confined in a Paul trap eventually reaches a Coulomb crystalline state when strongly cooled toward absolute zero. The normalized emittance of the Coulomb crystal can be in the sub-femtometer range. The trap is thus usable as a unique ion source for nano-beam production, though the available beam intensity is limited. This new concept was first discussed nearly 20 years ago *and later experimentally demonstrated by several research groups* (\*\*). In this paper, we report on the result of a recent experiment where an attempt was made to extract Ca<sup>+</sup> or N<sup>2+</sup> ions one by one from a compact linear Paul trap. In addition to the regular extraction scheme based on a string Coulomb crystal, the possibility of using a multi-shell crystalline structure is explored in detail.

### Footnotes

- M. Kano et al., J. Phys. Soc. Jpn. 73, 760(2004). \*\* W. Schnitzler et al., Phys. Rev. Lett. 102, 070501 (2009). \*\*\* K. Izawa et al., J. Phys. Soc. Jpn. 79, 124502 (2010).

### Funding Agency

### Paper preparation format

LaTeX

### Region represented

Asia

**Primary author:** MUROO, Kento (Hiroshima University)

**Co-authors:** ITO, Kiyokazu (Hiroshima University); OKAMOTO, Hiromi (Hiroshima University)

**Presenter:** MUROO, Kento (Hiroshima University)

**Session Classification:** Student Poster Session

**Track Classification:** MC3: Novel Particle Sources and Acceleration Techniques: MC3.T01 Proton and Ion Sources