



Contribution ID: 2252 Contribution code: SUPG074

Type: **Poster Presentation**

## Investigation of hot-spot and quench location due to trapped flux in niobium superconducting radiofrequency cavities

*Sunday, 19 May 2024 16:00 (2 hours)*

One of the significant sources of residual losses in superconducting radio-frequency cavities is magnetic flux trapped during the cool-down due to the incomplete Messier effect. If the trapped vortices are non-uniformly distributed on the cavity surface, the temperature mapping revealed the “hot-spots” at the location of high density of pinned vortices. In this contribution, we present the results of combined temperature and magnetic mapping measurements on a single cell 3.0 GHz and 1.3 GHz niobium single-cell cavities. The results show the direct evidence of pinned vortices induced “hot-spots”.

### Footnotes

### Funding Agency

The work is partially supported by the U.S. DOE, Office of science, Office of High Energy Physics under Awards No. DE-SC 0009960. This manuscript has been authored by Jefferson Science Associates, L

### Paper preparation format

### Region represented

Asia

**Primary author:** KHANAL, Bashu (Old Dominion University)

**Co-authors:** CIOVATI, Gianluigi (Old Dominion University); DHAKAL, Pashupati (Thomas Jefferson National Accelerator Facility)

**Presenter:** KHANAL, Bashu (Old Dominion University)

**Session Classification:** Student Poster Session

**Track Classification:** MC7: Accelerator Technology and Sustainability: MC7.T07 Superconducting RF